

AD-A223 658

DTIC FILE COPY

2

CHEMICAL  
RESEARCH,  
- DEVELOPMENT &  
ENGINEERING  
CENTER

CRDEC-TR-155

TRIMETHYLTIN SELECTIVELY ALTERS  
ACTIVITY OF  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$ , AND  
( $\text{Ca}^{++}$  +  $\text{Mg}^{++}$ )-ATPases OF  
HUMAN NEUROBLASTOMA

James J. Valdes, Ph.D.  
Roy G. Thompson  
RESEARCH DIRECTORATE

James P. Chambers, Ph.D.  
UNIVERSITY OF TEXAS AT SAN ANTONIO  
San Antonio, TX 78285

May 1990

DTIC  
ELECTE  
JUN 28 1990  
S E D

U.S. ARMY  
ARMAMENT  
MUNITIONS  
CHEMICAL COMMAND



Aberdeen Proving Ground, Maryland 21010-8423

Best Available Copy

00 00 00 008

#### Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.

#### Distribution Statement

Approved for public release; distribution is unlimited.

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20543.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 1990 May		3. REPORT TYPE AND DATES COVERED Final, 86 Oct - 86 Nov	
4. TITLE AND SUBTITLE Trimethyltin Selectively Alters Activity of $Ca^{++}$ , $Mg^{++}$ , and $(Ca^{++} + Mg^{++})$ -ATPases of Human Neuroblastoma				5. FUNDING NUMBERS PR-1C162706A553	
6. AUTHOR(S) Valdes, James J., Ph.D., and Thompson, Roy G. (CRDEC); and Chambers, James P., Ph.D. (University of Texas at San Antonio)					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Texas at San Antonio San Antonio, TX 78285				8. PERFORMING ORGANIZATION REPORT NUMBER CRDEC-TR-155	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Cdr, CRDEC, ATTN: SMCCR-RSB, APG, MD 21010-5423				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Human neuroblastoma GM3320 tissue homogenates exhibit $Ca^{++}$ , $Mg^{++}$ , and $(Ca^{++} + Mg^{++})$ -ATPase activities which are ouabain insensitive. Trimethyltin (TMT) discriminates among these three ATPase activities. At low (25 $\mu$ M) concentrations, TMT stimulated the $Ca^{++}$ -ATPase and inhibited the $(Ca^{++} + Mg^{++})$ -ATPase by 70%. At 75 $\mu$ M, TMT inhibited the $Mg^{++}$ -ATPase by 50% and the $(Ca^{++} + Mg^{++})$ -ATPase by 95%. In summary, TMT preferentially inhibited the $Mg^{++}$ and $(Ca^{++} + Mg^{++})$ -ATPase activity of human neuroblastoma GM3320 cells.					
14. SUBJECT TERMS Trimethyltin      Neuroblastoma Organotin ATPase				15. NUMBER OF PAGES 13	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL		

Blank

# PREFACE

The work described in this report was authorized under Project No. 1C162706A553C, Decontamination, Detection and Identification. This work was started in October 1986 and completed in November 1986. The experimental data are contained in laboratory notebooks in the Division of Life Sciences, University of Texas at San Antonio (San Antonio, TX).

The use of trade names or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, U.S. Army Chemical Research, Development and Engineering Center, ATTN: SMCCR-SPS-T, Aberdeen Proving Ground, Maryland 21010-5423. However, the Defense Technical Information Center and the National Technical Information Service are authorized to reproduce this document for U.S. Government purposes.

This report has been approved for release to the public.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Special
A-1	



Blank

## CONTENTS

	Page
1. INTRODUCTION .....	7
2. MATERIALS AND METHODS .....	7
2.1 Materials .....	7
2.2 Cell Culture .....	8
2.3 Enzyme Assays .....	8
3. RESULTS .....	8
4. DISCUSSION .....	10
5. CONCLUSIONS .....	10
LITERATURE CITED .....	11

Blank



# TRIMETHYLTIN SELECTIVELY ALTERS ACTIVITY OF $\text{Ca}^{++}$ , $\text{Mg}^{++}$ , AND $(\text{Ca}^{++} + \text{Mg}^{++})$ -ATPases OF HUMAN NEUROBLASTOMA

## 1. INTRODUCTION

Neurotransmitter release from nerve terminals is coupled to the intracellular concentration of  $\text{Ca}^{++}$ . The regulation of internal  $\text{Ca}^{++}$  pools is, in turn, accomplished by several processes which include the transport of  $\text{Ca}^{++}$  into mitochondria,<sup>1</sup> the binding of  $\text{Ca}^{++}$  by intraneuronal or membrane proteins,<sup>2,3</sup> the exchange of intraneuronal  $\text{Ca}^{++}$  with extracellular  $\text{Na}^{+}$  via an antiport process,<sup>4,5</sup> and finally,  $\text{Ca}^{++}$  extrusion or sequestration by  $\text{Ca}^{++}$ -activated adenosine triphosphatases (ATPases).<sup>6-9</sup> These processes are sensitive targets for a wide range of neurotoxic compounds with industrial, agricultural, and military applications. In fact,  $(\text{Ca}^{++} + \text{Mg}^{++})$ -ATPase and an associated ion channel are inhibited by Mojave toxin<sup>10</sup> and activated by maitotoxin,<sup>11</sup> two of the most potent known naturally occurring neurotoxins. (JL)<sup>6</sup>

Organotin compounds are biologically active organometals which are used as pesticides, fungicides, anti-fouling paints, plastic stabilizers, and catalysts.<sup>12,13</sup> The short chain alkyltins are water soluble and degrade to relatively non-toxic products, but induce neuropathies in mammals.<sup>14-19</sup> Trimethyltin (TMT), the most potent neurotoxic organotin compound,<sup>20</sup> induces psychological and behavioral disturbances in humans characterized by irritability, memory loss, and mental confusion.<sup>21-23</sup>

Neurochemical effects include alterations of muscarinic and dopaminergic receptor binding in mice,<sup>24</sup> inhibition of neurotransmitter uptake in mouse<sup>25</sup> and rat<sup>19</sup> brain synaptosomes, and inhibition of acetylcholine release under conditions of 20 Hz stimulation in the rat phrenic nerve-hemidiaphragm.<sup>26</sup> The apparent nonspecificity of TMT for neurotransmitter systems and the similarity of its effects to those of ruthenium red, which blocks  $\text{Ca}^{++}$  entry into mitochondria,<sup>27</sup> suggest that TMT may disturb membrane pumps responsible for the regulation of intracellular  $\text{Ca}^{++}$ . These observations and the fact that triethyltin, a related compound, has general effects on enzymes which use ATP as a substrate,<sup>28-30</sup> provide the rationale for assessing the effects of TMT on endogenous ATPase activities in human neuroblastoma cells.

## 2. MATERIALS AND METHODS

### 2.1 Materials.

Trimethyltin chloride and malachite green were purchased from Aldrich Chemical Company, Milwaukee, Wisconsin. Adenosine triphosphate (ATP), ouabain, HEPES buffer, Tris buffer, and glycerol were purchased from Sigma Chemical Company, St. Louis, Missouri. Dulbecco Modified Eagle's media (DMEM-1X) was purchased from Irvine Scientific, Irvine, California. Fetal Calf Serum (FCS) was purchased from Hyclone Labs, Logan, Utah. Trypsin was obtained from Gibco, Grand Island, Maine. Human neuroblastoma cells (GM3320) were obtained from Coriell Human Genetic Mutant Cell Repository, Camden, New Jersey.

## 2.2 Cell Culture.

Cells were grown at 37°C in Falcon T-75 flasks containing 10 ml DME-1X supplemented with 10% (v/v) HCS in an atmosphere of 1% CO<sub>2</sub>. Media was changed every three days. Confluency, approximately  $1.0 \times 10^{-7}$  cells, was attained on the 7th or 8th day, at which time the culture flasks were treated for 1 min with a 0.05% trypsin solution containing 0.05% EDTA (w/v). Trypsin action was inhibited by addition of an equal volume of growth medium, followed immediately by centrifugation for 5 min (1500 rpm) at room temperature.

The resulting pellet was washed twice with 0.01 M Tris buffer, pH 8.2, containing 16% (v/v) glycerol. The washed pellet was then homogenized (10 strokes) in a Potter Elvehjem glass homogenizer previously chilled to 5°C. The protein was determined by the method of Bradford using bovine serum albumin as a standard.<sup>31</sup>

## 2.3 Enzyme Assays.

All incubations had a final volume of 2.0 ml containing 150 µg protein, 0.02 M HEPES buffer, pH 7.2, 100 µM EGTA, 100 mM KCl, 0.1 mM ouabain, indicated amounts of TMT, and 250 µM ATP. Prior to the addition of ATP, assay mixtures were preincubated for 5 min at 37°C. Hydrolysis of ATP was allowed to proceed for 10 min at 37°C. Reactions were terminated by addition of 200 µl 6.0 N HCl at which time 200 µl aliquots were removed. Released phosphate was monitored spectrophotometrically by the method of Lanzetta and coworkers.<sup>32</sup> Ca<sup>++</sup>-ATPase activity was determined by subtracting basal ATPase activity (no added Ca<sup>++</sup> or Mg<sup>++</sup>) from activity in the presence of 300 µM CaCl<sub>2</sub> (no Mg<sup>++</sup>). Mg<sup>++</sup>-ATPase activity was the difference between basal and Mg<sup>++</sup>-stimulated (1.0 mM) activity (no Ca<sup>++</sup>). (Ca<sup>++</sup> + Mg<sup>++</sup>)-dependent ATPase activity represents the difference between total activity (Ca<sup>++</sup> + Mg<sup>++</sup>) and activity in the presence of Mg<sup>++</sup> alone. Assay conditions described above were linear with respect to both protein and time. Trimethyltin chloride solutions were prepared fresh in glass distilled water: 25, 50, 75, 100, and 125 µM final concentrations.

## 3. RESULTS

As shown in the figure, very low concentrations of TMT significantly depress the Ca<sup>++</sup>, Mg<sup>++</sup>, and (Ca<sup>++</sup> + Mg<sup>++</sup>)-dependent ATPase activities of neuroblastoma GM3320 homogenates. At 25 µM, the (Ca<sup>++</sup> + Mg<sup>++</sup>)-dependent ATPase activity (●-●-●) is inhibited by greater than 65% in contrast to a 45% stimulation of the Ca<sup>++</sup>-dependent, Mg<sup>++</sup>-independent activity (o-o-o); whereas, the Mg<sup>++</sup>-dependent, Ca<sup>++</sup>-independent ATPase activity (x-x-x) was observed to be inhibited by approximately 25%. At 75 µM, no inhibition of the Ca<sup>++</sup>-dependent, Mg<sup>++</sup>-independent ATPase activity was observed; whereas, the (Ca<sup>++</sup> + Mg<sup>++</sup>)-dependent activity was inhibited greater than 95%. Only a moderate inhibition of approximately 40% of the Mg<sup>++</sup>-dependent, Ca<sup>++</sup>-independent activity was observed at 75 µM TMT. Concentration of greater than 75 µM TMT in the respective assays resulted in increasing inhibition of all activities.




Figure. The effect of increasing amounts of triethyltin (TMT) on Ca<sup>++</sup>, Mg<sup>++</sup>, and (Ca<sup>++</sup> + Mg<sup>++</sup>)-dependent ATPase activities of an neuroblastoma GM320. Ca<sup>++</sup>-dependent, Mg<sup>++</sup>-independent ATPase activity (o-o-o); Mg<sup>++</sup>-dependent, Ca<sup>++</sup>-independent ATPase activity (x-x-x); and (Ca<sup>++</sup> + Mg<sup>++</sup>)-dependent ATPase activity (●-●-●).

#### 4. DISCUSSION

At present, little is known about the mechanism of the neurotoxicity of TMT. A related compound, triethyltin, has been reported to inhibit oxidative phosphorylation and to decrease the incorporation of ( $^{14}\text{C}$ )-glucose into pyruvate and several putative amino acid transmitters.<sup>21</sup> This might account for the observed toxicity to cultured neuroblastoma GB320 cells of very low concentrations of TMT. Data presented here indicate that TMT may be useful as an activator or inhibitor of ATPase activity in disrupted tissue homogenates. This is consistent with the idea that alkyltins, such as TMT, have a high affinity for membranes and processes associated with membranes as in the inhibition by TMT of the uptake of neurotransmitters into synaptosomes. These results support the contention that this compound may be useful for studying ATP dependent processes such as neurotransmitter release. Recent reports<sup>33,34</sup> suggest that the  $\text{Ca}^{++}$ -ATPase,  $\text{Mg}^{++}$ -ATPase and  $(\text{Ca}^{++} + \text{Mg}^{++})$ -ATPase activities in synaptic plasma membranes reflect the operation of three separate enzymes. Although these activities must be studied in greater detail, their presence suggest the potential usefulness of this cell line in the study of various ATPase dependent neurochemical processes.

#### 5. CONCLUSIONS

The  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$ , and  $(\text{Ca}^{++} + \text{Mg}^{++})$ -ATPases are selectively sensitive to perturbation by TMT, suggesting a role for these enzymes in the neuropathologies induced by this organotin. The neuroblastoma GB320 cell culture provides a convenient system with which to study the mechanisms of action of a wide variety of neurotoxic agents and to assess the potential of these enzymes as detectors of neurotoxins.

# LITERATURE CITED

1. Blaustein, M.P., Ratzlaff, R.W., and Kendrick, N.K. "The Regulation of Intracellular Calcium in Presynaptic Nerve Terminals," Ann. N.Y. Acad. Sci. Vol. 307, pp 195-212 (1978).
2. Cheung, W.Y. "Cyclic 3', 5'-Nucleotide Phosphodiesterase," J. Biol. Chem. Vol. 246, pp 2859-2869 (1971).
3. Teo, T.S. and Wang, J.H. "Mechanism of Activation of a Cyclic Adenosine 3':5'-monophosphate Phosphodiesterase from Bovine Heart by Calcium Ions," J. Biol. Chem. Vol. 248, pp 5950-5955 (1973).
4. Blaustein, M.P. and Oborn, C.J. "The Influence of Sodium on Calcium Fluxes in Pinched-off Nerve Terminals in Vitro," J. Physiol. (Lond.) Vol. 247, pp 657-686 (1975).
5. Michaelis, M.L. and Michaelis, E.K. "Ca<sup>++</sup> Fluxes in Released Synaptic Plasma Membrane Vesicles," Life Sci. Vol. 28, pp 37-45 (1981).
6. Javors, M.A., Bowden, C.L., and Ross, D.H. "Kinetic Characterization of Ca<sup>++</sup> Transport in Synaptic Membranes," J. Neurochem. Vol. 37, pp 381-387 (1981).
7. Kuo, C.-H., Ichida, S., Matsuda, T., Kakiuchi, S., and Yoshida, H. "Regulation of ATP-dependent Ca<sup>++</sup> Uptake of Synaptic Plasma Membranes by Ca<sup>++</sup> Dependent Modular Protein," Life Sci. Vol. 25, pp 235-240 (1979).
8. Rahamimoff, H. and Abramovitz, E. "Calcium Transport in a Vesicular Membrane Preparation from Rat Brain Synaptosomes," FEBS. Lett. Vol. 89, pp 223-226 (1978).
9. Rahamimoff, H. and Abramovitz, E. "Calcium Transport and ATPase Activity of Synaptosomal Vesicles from Rat Brain," FEBS Lett. Vol. 92, pp 163-167 (1978).
10. Chambers, J.P., Wayner, M.J., Dungan, J., Rael, E.D., and Valdes, J.J. "The Effects of Purified Mojave Toxin on Rat Synaptic Membrane (Ca<sup>++</sup>-Mg<sup>++</sup>)-ATPase and the Dihydropyridine Receptor," Brain Res. Bull. Vol. 16, pp 639-643 (1986).
11. Takahashi, M., Tatsumi, M., and Ohizumi, Y. "Ca<sup>++</sup> Channel Activating Function of Maitotoxin, the Most Potent Marine Toxin Known, in Clonal Rat Pheochromocytoma Cells," J. Biol. Chem. Vol. 258(18), pp 10944-10948 (1983).
12. Piver, W.T. "Organotin Compounds: Industrial Applications and Biological Investigations," Environ. Health Perspect. Vol. 4, pp 61-79 (1973).
13. Hunter, R.C. "Organotin Compounds and Their Use for Insect and Mite Control," Environ. Health Perspect. Vol. 14, p 47 (1976).

14. Bouldin, T.W., Goines, N.D., Bagnell, C.R., and Krigman, M.R. "Pathogenesis of Trimethyltin Neuronal Toxicity," Am. J. Pathol. Vol. 104, pp 237-249 (1981).
15. Brown, A.W., Aldridge, W.N., Street, B.W., and Verschoyle, R.D. "The Behavior and Neuropathologic Sequelae of Intoxication by Trimethyltin Compounds in the Rat," Am. J. Pathol. Vol. 97, pp 59-76 (1979).
16. Dyer, R.S., Walsh, T.J., Wonderlin, W.F., and Bercegeay, M. "The Trimethyltin Syndrome in Rats," Neurobehav. Toxicol. Teratol. Vol. 4, pp 127-133 (1982).
17. Dyer, R.S., Deshields, T.L., and Wonderlin, W.F. "Trimethyltin-induced Changes in Gross Morphology of the Hippocampus," Neurobehav. Toxicol. Teratol. Vol. 4, pp 141-147 (1982).
18. Dyer, R.S., Wonderlin, W.F., and Walsh, T.J. "Increased Seizure Susceptibility Following Trimethyltin Administration in Rats," Neurobehav. Toxicol. Teratol. Vol. 4, pp 203-208 (1982).
19. Valdes, J.J., Mactutus, C.F., Santos-Anderson, R.M., Dawson, R., and Annau, Z. "Selective Neurochemical and Histological Lesions in Rat Hippocampus Following Chronic Trimethyltin Exposure," Neurobehav. Toxicol. Teratol. Vol. 5, pp 357-361 (1983).
20. Stoner, H.B., Earnes, J.M., and Duff, J.I. "Studies on the Toxicity of Alkyltin Compounds," Br. J. Pharmacol. Chemother. Vol. 10, pp 16-25 (1955).
21. Cremer, J.E. "Selective Inhibition of Glucose Oxidation by Triethyltin in Rat Brain in Vivo," Biochem. J. Vol. 119, pp 95-102 (1981).
22. Fortemps, E., Amand, G., Bomboir, A., Lauwerys, R., and Laterre, E.C. "Trimethyltin Poisoning-Report of Two Cases," Int. Arch. Occup. Environ. Health. Vol. 41, pp 1-6 (1978).
23. Ross, W.D., Emmett, E.A., Steiner, J., and Tureen, R. "Neurotoxic Effects of Occupational Exposure to Organotins," Am. J. Psych. Vol. 138, pp 1092-1103 (1981).
24. Ali, S.F., Cranmer, J.M., Goad, P.T., Slikker, W., Harbison, R.D., and Cranmer, M.F. "Trimethyltin Induced Changes of Neurotransmitter Levels and Brain Receptor Binding in the Mouse," Neurotoxicology Vol 4, pp 29-36 (1983).
25. Doctor, S.V., Costa, L.G., Kendall, D.A., and Murphy, S.D. "Trimethyltin Inhibits Uptake of Neurotransmitters into Mouse Forebrain Synaptosomes," Toxicology Vol. 25, pp 213-221 (1982).
26. Bierkamper, G.G. and Valdes, J.J. "Triethyltin Intoxication Alters Acetylcholine Release from Rat Phrenic Nerve-hemidiaphragm," Neurobehav. Toxicol. Teratol. Vol. 4, pp 251-254 (1982).

27. Rahaminoff, R. "The Regulation of Intracellular Calcium Concentration and Transmitter Release," Neurosci. Res. Prog. Bull. Vol. 15, pp 575-581 (1979).
28. Davidoff, F. and Carr, S. "Interaction of Triethyltin with Pyruvate Kinase," Biochemistry Vol. 12, pp 1415-1422 (1973).
29. Siebenlist, K.R. and Takeda, F. "Inhibition of Red Cell and Yeast Hexokinase by Triethyltin Bromide," Biochem. Biophys. Res. Commun. Vol. 95, pp 758-764 (1980).
30. Siebenlist, K.R. and Takeda, F. "The Effects of Triethyltin Bromide on Red Cell and Brain Cyclic AMP-dependent Protein Kinases," J. Biol. Chem. Vol. 258, pp 11384-11390 (1983).
31. Brauford, M. "A Rapid and Sensitive Method for the Quantification of Microgram Quantities of Protein Utilizing the Principle of Protein-dye Binding," Anal. Biochem. Vol. 72, pp 248-254 (1976).
32. Lanzetta, P., Alvarez, P., Reinach, P., and Candia, O. "An Improved Assay for Nanomole Amounts of Inorganic Phosphate," Anal. Biochem. Vol. 100, pp 95-97 (1979).
33. Lin, S. and Way, E.L. "A High Affinity  $\text{Ca}^{++}$  ATPase in Enriched Nerve-ending Plasma Membranes," Brain Research Vol. 235, pp 387-392 (1982).
34. Papazian, D.M., Rahaminoff, H., and Goldin, S.M. "Partial Purification and Functional Identification of a Calmodulin-activated, Adenosine 5'-triphosphate-dependent Calcium Pump from Synaptic Plasma Membranes," J. Neurosci. Vol. 4, pp 1933-1943 (1984).